



(continued)

19. Confirm initial screen was accomplished, verify results, and follow up, as appropriate. The Recommended Uniform Screening Panel (<https://www.hrsa.gov/advisory-committees/heritable-diseases/rusp/index.html>), as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<http://gensr-usa.uihscs.edu/home>) establish the criteria for and coverage of newborn screening procedures and programs.
20. Verify results as soon as possible, and follow up, as appropriate.
21. Confirm initial screening was accomplished, verify results, and follow up, as appropriate. See "Hyperbilirubinemia in the Newborn Infant: 35 Weeks of Age With Clinicians" (<http://pediatrics.aappublications.org/content/124/4/1193>).
22. Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per "Encouragement of Prenatal and Neonatal Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<http://pediatrics.aappublications.org/content/129/7/1906>).
23. Schedules, per the AAP Committee on Infectious Diseases, are available at [http://redbook.solutions.aap.org/SS/immunization\\_Schedules.aspx](http://redbook.solutions.aap.org/SS/immunization_Schedules.aspx). Every visit should be an opportunity to update and complete a child's immunizations.
24. Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP *Pediatric Nutrition: Policy of the American Academy of Pediatrics* (Iron chapter).
25. For children at risk of lead exposure, see "Prevention of Childhood Lead Toxicity" (<http://pediatrics.aappublications.org/content/138/7/e2016.1493>) and "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" ([http://www.cdc.gov/nceh/lead/ACCLPP/Final\\_Document\\_03/07/2.pdf](http://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_03/07/2.pdf)).
26. Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.
27. Tuberculosis testing per recommendations of the AAP Committee on Infectious Diseases, published in the current edition of the *AAP Red Book: Report of the Committee on Infectious Diseases*. Testing should be performed on recognition of high-risk factors.
28. See "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" ([http://www.heart.org/healthguidelines/child\\_ped/index.html](http://www.heart.org/healthguidelines/child_ped/index.html)).
29. Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the *AAP Red Book: Report of the Committee on Infectious Diseases*.
30. Adolescents should be screened for HIV according to the USPSTF recommendations between the ages of 15 and 18, making every effort to preserve confidentiality of the adolescent. Those at increased risk of HIV infection, including those who are sexually active, participate in injection drug use, or are being tested for other STIs, should be tested for HIV and reassessed annually.
31. See USPSTF recommendations (<https://www.uspreventiveservicestaskforce.org/PracticeGuidance/updates/unananimous/neck-cancer-screening>). Indications for pelvic examinations prior to age 21 are noted in "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<http://pediatrics.aappublications.org/content/126/3/583>, full).
32. Assess whether the child has a dental home. If no dental home is identified, perform a risk assessment (<https://www.aap.org/on-us/infocenter-and-policies/aap-health-initiatives/Oral-Health/Oral-Health-Practice-Tool.aspx>) and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See "Maintaining and Improving the Oral Health of Young Children" (<http://pediatrics.aappublications.org/content/134/6/11224>).
33. Perform a risk assessment. See "Maintaining and Improving the Oral Health of Young Children" (<http://pediatrics.aappublications.org/content/134/6/11224>).
34. See USPSTF recommendations (<https://www.uspreventiveservicestaskforce.org/PracticeGuidance/updates/unanimous/fluoride-toothpastes>). The recommendations should be applied to all children every 3-6 months in the primary care or dental office. Indications for fluoride use are noted in "Fluoride Use in Caries Prevention in the Primary Care Setting" (<http://pediatrics.aappublications.org/content/134/3/629>).
35. If primary water source is deficient in fluoride, consider oral fluoride supplementation. See "Fluoride Use in Caries Prevention in the Primary Care Setting" (<http://pediatrics.aappublications.org/content/134/3/629>).

## Bright Futures/AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule)

This schedule reflects changes approved in December 2018 and published in March 2019. For updates and a list of previous changes made, visit [www.aap.org/periodicity/schedule](http://www.aap.org/periodicity/schedule).

### CHANGES MADE IN DECEMBER 2018

#### BLOOD PRESSURE

- Footnote 6 has been updated to read as follows: "Screening should occur per Clinical Practice Guideline for Screening and Management of High Blood Pressure in Children and Adolescents" (<http://pediatrics.aappublications.org/content/140/3/e20171904>). Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years."

#### ANEMIA

- Footnote 24 has been updated to read as follows: "Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP *Pediatric Nutrition: Policy of the American Academy of Pediatrics* (Iron chapter)."

#### LEAD

- Footnote 25 has been updated to read as follows: "For children at risk of lead exposure, see 'Prevention of Childhood Lead Toxicity' (<http://pediatrics.aappublications.org/content/138/7/e2016.1493>) and 'Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention' ([https://www.cdc.gov/nceh/lead/ACCLPP/Final\\_Document\\_03/07/2.pdf](https://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_03/07/2.pdf))."



This program is supported by the Health Resources and Services Administration, U.S. Department of Health and Human Services (HRSA) as part of an award totaling \$5,000,000 with 10 percent financed with non-governmental sources. The contents are those of the author(s) and do not necessarily represent the official views or recommendations of HRSA, HHS, or the U.S. Government. For more information, please visit [HRSA.gov](http://HRSA.gov).