

Immunization Timing 2024

Between 0-6 months, one dose of RSV vaccine may be needed for some babies.



At Birth

HepB
(Hepatitis B)

Age 0-2 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

HepB
(Hepatitis B)
1-2 months

Hib
(Hib meningitis)

PCV
(Pneumo)

RV
(Rotavirus)

Age 4 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

Hib
(Hib meningitis)

PCV
(Pneumo)

RV
(Rotavirus)

Age 6 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

HepB
(Hepatitis B)

PCV
(Pneumo)

RV
(Rotavirus)

Age 12 months

HepA
(Hepatitis A)

MMR
(measles, mumps, and rubella)

Hib
(Hib meningitis)

PCV
(Pneumo)

Varicella
(Chicken Pox)

Age 15 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Age 18 months

HepA
(Hepatitis A)

Age 4-6 years	Age 11-12 years	Age 16-18 years
DTaP Polio (IPV) MMR Varicella	Tdap HPV (2 doses) MenACWY (Meningitis)	MenACWY (Meningitis) MenB

Please ask your doctor how far apart to have the vaccines. For more information, visit: www.cdc.gov/vaccines.

COVID-19 vaccination is recommended for ages 6 months and older.

Keep your child safe from the flu. Everyone 6 months and older should get the flu vaccine. To best protect your child, get them the flu vaccine during the fall season.



CALIFORNIA KIDS
Love them. Immunize them.