



# Start Well Childhood Vaccines

## IMMUNIZATION TIMING 2026

Kids need vaccines to protect them from serious diseases.

Within the first year, some babies may need one dose of RSV. Talk to your doctor to see what's right for your baby.

COVID-19 vaccination is recommended for ages 6 months and older.

To best protect your child, get them a flu vaccine every fall. Everyone 6 months and older should get the flu vaccine. To best protect your child, get them the flu vaccine during the fall season.



			<b>Age 6 months</b>	
	<b>Age 2 months</b>		<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	
	<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	<b>Age 4 months</b>	<b>IPV</b> (Polio)	<b>Age 12 months</b>
	<b>IPV</b> (Polio)	<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	<b>HepB</b> (Hepatitis B)	<b>HepA</b> (Hepatitis A)
	<b>HepB</b> (Hepatitis B)	<b>IPV</b> (Polio)	<b>Hib</b> (Hib meningitis)	<b>MMR</b> (Measles, Mumps and Rubella)
	<b>Hib</b> (Hib meningitis)	<b>Hib</b> (Hib meningitis)	<b>PCV</b> (Pneumo)	<b>Hib</b> (Hib meningitis)
	<b>PCV</b> (Pneumo)	<b>PCV</b> (Pneumo)	<b>RV</b> (Rotavirus)	<b>PCV</b> (Pneumo)
	<b>HepB</b> (Hepatitis B)			<b>Age 15 months</b>
	<b>RV</b> (Rotavirus)	<b>RV</b> (Rotavirus)	<b>Flu</b> 2 shots one month apart	<b>Age 18 months</b>
				<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)
				<b>HepA</b> (Hepatitis A)

**Age 4-6 YEARS**

**DTaP**  
**Polio**  
**MMR**  
**Varicella**

**Age 11-12 YEARS**

**Tdap**  
**HPV**  
(2 doses)  
**Meningitis**

**Age 16-18 YEARS**

**Meningitis**  
**Meningitis B**  
(2 doses)

Please ask your doctor how far apart to have the vaccines. For more information, visit: [AAP.org/ImmunizationSchedule](https://www.aap.org/ImmunizationSchedule)