

Immunization Timing 2023

At Birth

HepB
(Hepatitis B)

Age 0-2 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

HepB
(Hepatitis B) 1-2 months

Hib
(Hib meningitis)

PCV
(Pneumo)

RV
(Rotavirus)

Age 4 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

Hib
(Hib meningitis)

PCV
(Pneumo)

RV
(Rotavirus)

Age 6 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Polio
(IPV)

HepB
(Hepatitis B)

PCV
(Pneumo)

RV
(Rotavirus)

Age 12 months

HepA
(Hepatitis A)

MMR
(measles, mumps, and rubella)

Hib
(Hib meningitis)

PCV
(Pneumo)

Varicella
(Chicken Pox)

COVID-19 vaccination is recommended for ages 6 months and older.



| Age 4-6 years | Age 11-12 years | Age 16-18 years |
|---|---|------------------------------|
| DTaP Polio (IPV) MMR Varicella | Tdap HPV (2 doses) MenACWY (Meningitis) | MenACWY (Meningitis) MenB |

Age 15 months

DTaP
(Diphtheria, Tetanus, Pertussis)

Age 18 months

HepA
(Hepatitis A)

Please ask your Doctor how far apart to have the vaccines. For more information, visit: www.cdc.gov/vaccines

California schools are requiring COVID-19 vaccines for some kids. The requirement is subject to change. Please go to www.ShotsForSchool.org for the most current information.

Flu Vaccine Every Fall

Everyone 6 months and older should get one.



CALIFORNIA KIDS
Love them. Immunize them.